arXiv:math/0010223v1 [math.GT] 24 Oct 2000

HOMOTOPY TYPES OF HOMEOMORPHISM GROUPS OF NONCOMPACT 2-MANIFOLDS

TATSUHIKO YAGASAKI

Abstract. Suppose M is a noncompact connected PL 2-manifold and let $\mathcal{H}(M)_0$ denote the identity component of the homeomorphism group of M with the compact-open topology. In this paper we classify the homotopy type of $\mathcal{H}(M)_0$ by showing that $\mathcal{H}(M)_0$ has the homotopy type of the circle if M is the plane, an open or half open annulus, or the punctured projective plane. In all other cases we show that $\mathcal{H}(M)_0$ is homotopically trivial.

1. Introduction

Hamstrom [4] classified the homotopy types of the identity components of homeomorphism groups of compact 2-manifolds M. In this paper we treat the case where M is noncompact. Suppose M is a PL 2-manifold and X is a compact subpolyhedron of M. We denote by $\mathcal{H}_X(M)$ the group of homeomorphisms h of M onto itself with $h|_X = id$, equipped with the compact-open topology, and by $\mathcal{H}(M)_0$ the identity component of $\mathcal{H}(M)$. Let $\mathbb{R}^2$ denote the plane, $\mathbb{S}^1$ the unit circle and $\mathbb{P}^2$ the projective plane. The following is the main result of this paper.

Theorem 1.1. *Suppose M is a noncompact connected (separable) PL 2-manifold and X is a compact subpolyhedron of M. Then*
(i) $\mathcal{H}_X(M)_0 \simeq \mathbb{S}^1$ *if* $(M, X) \cong (\mathbb{R}^2, \emptyset)$, $(\mathbb{R}^2, 1pt)$, $(\mathbb{S}^1 \times \mathbb{R}^1, \emptyset)$, $(\mathbb{S}^1 \times [0,1), \emptyset)$ *or* $(\mathbb{P}^2 \setminus 1pt, \emptyset)$,
(ii) $\mathcal{H}_X(M)_0 \simeq *$ *in all other cases.*

Corollary 1.1. *If M is a connected (separable) 2-manifold and X is a compact subpolyhedron of M with respect to some triangulation of M, then $\mathcal{H}_X(M)_0$ is an ℓ_2-manifold.*

In [14] we obtained a natural principal bundle connecting the homeomorphism group and the embeddding space (cf. Section 2). In this paper we will seek a condition under which the fiber of this bundle is connected (Section 3). The contractibility and the ANR property of $\mathcal{H}_X(M)_0$ in the compact case will then imply the similar properties of embedding spaces and in turn the corresponding properties of $\mathcal{H}_X(M)_0$ in the noncompact case. Corollary 1.1 follows immediately from the characterization of ℓ_2-manifolds and this enables us to determine the topological type itself of $\mathcal{H}_X(M)_0$ by the homotopy invariance of infinite-dimensional manifolds.

In a succeeding paper we will investigate the subgroups of $\mathcal{H}_X(M)_0$ consisting of PL and Lipschitz homeomorphisms from the viewpoints of infinite-dimensional topological manifolds.

1991 *Mathematics Subject Classification.* 57N05, 57N20.

Key words and phrases. 2-manifolds, Homeomorphism groups, ℓ_2-manifolds.

2. Preliminaries

Throughout the paper we follow the following conventions: Spaces are assumed to be separable and metrizable, and maps are always continuous. When A is a subset of a space X, the notations $\mathrm{Fr}_X A$, $\mathrm{cl}_X A$ and $\mathrm{Int}_X A$ denote the frontier, closure and interior of A relative to X (i.e., $\mathrm{Int}_X A = \{x \in A \mid A$ contains a neighborhood of x in $X\}$ and $\mathrm{Fr}_X A = \mathrm{cl}_X A \setminus \mathrm{Int}_X A$). On the other hand, when M is a manifold, the notations ∂M and $\mathrm{Int}\, M$ denote the boundary and interior of M as a manifold. When N is a 2-submanifold of a 2-manifold M, we always assume that N is a closed subset of M and $\mathrm{Fr} N = \mathrm{Fr}_M N$ is a 1-manifold transversal to ∂M. Therefore we have $\mathrm{Int}\, N = \mathrm{Int}_M N \cap \mathrm{Int}\, M$ and $\mathrm{Fr}_M N \subset \partial N$. A metrizable space X is called an ANR (absolute neighborhood retract) if any map $f : B \to X$ from a closed subset of a metrizable space Y has an extension to a neighborhood U of B. If we can always take $U = Y$, then X is called an AR (absolute retract). ANRs are locally contractible and ARs are exactly contractible ANRs (cf. [7]). Finally ℓ_2 denotes the separable Hilbert space $\{(x_n) \in \mathbb{R}^\infty : \sum_n x_n^2 < \infty\}$.

In [14] we investigated some extension property of embeddings of a compact 2-polyhedron into a 2-manifold, based upon the conformal mapping theorem. The result is summarized as follows: Suppose M is a PL 2-manifold and $K \subset X$ are compact subpolyhedra of M. Let $\mathcal{E}_K(X, M)$ denote the space of embeddings $f : X \hookrightarrow M$ with $f|_K = id$, equipped with the compact-open topology. We consider the subspace of proper embeddings $\mathcal{E}_K(X, M)^* = \{f \in \mathcal{E}_K(X, M) : f(X \cap \partial M) \subset \partial M, f(X \cap \mathrm{Int}\, M) \subset \mathrm{Int}\, M\}$. Let $\mathcal{E}_K(X, M)^*_0$ denote the connected component of the inclusion $i_X : X \subset M$ in $\mathcal{E}_K(X, M)^*$.

Theorem 2.1. *For every $f \in \mathcal{E}_K(X, M)^*$ and every neighborhood U of $f(X)$ in M, there exists a neighborhood $\mathcal{U}$ of f in $\mathcal{E}_K(X, M)^*$ and a map $\varphi : \mathcal{U} \to \mathcal{H}_{K \cup (M \setminus U)}(M)_0$ such that $\varphi(g) f = g$ for each $g \in \mathcal{U}$ and $\varphi(f) = id_M$.*

Corollary 2.1. *For any open neighborhood U of X in M, the restriction map $\pi : \mathcal{H}_{K \cup (M \setminus U)}(M)_0 \to \mathcal{E}_K(X, U)^*_0$, $\pi(f) = f|_X$, is a principal bundle with fiber $\mathcal{G} \equiv \mathcal{H}_{K \cup (M \setminus U)}(M)_0 \cap \mathcal{H}_X(M)$, where the group $\mathcal{G}$ acts on $\mathcal{H}_{K \cup (M \setminus U)}(M)_0$ by right composition.*

Proposition 2.1. *$\mathcal{E}_K(X, M)$ and $\mathcal{E}_K(X, M)^*$ are ANRs.*

Next we recall some fundamental facts on homeomorphism groups of compact 2-manifolds.

Fact 2.1. If N is a compact PL 2-manifold and Y is a compact subpolyhedron of N, then $\mathcal{H}_Y(N)$ is an ANR. ([8], [9], cf. [14, Lemma 3.2])

Lemma 2.1. ([4], [12, §3]) *Suppose N is a compact connected PL 2-manifold and Y is a compact subpolyhedron of N.*

(i) *If $(N, Y) \not\cong (\mathbb{D}^2, \emptyset)$, $(\mathbb{D}^2, 0)$, $(\mathbb{S}^1 \times [0,1], \emptyset)$, $(\mathbb{M}, \emptyset)$, $(\mathbb{S}^2, \emptyset)$, $(\mathbb{S}^2, 1pt)$, $(\mathbb{S}^2, 2pts)$, $(\mathbb{T}^2, \emptyset)$, $(\mathbb{K}^2, \emptyset)$, $(\mathbb{P}^2, \emptyset)$, $(\mathbb{P}^2, 1pt)$, then $\mathcal{H}_Y(N)_0 \simeq *$.*

(ii) *If A is a nonempty compact subset of ∂N, then $\mathcal{H}_{Y \cup A}(N)_0 \simeq *$.*

(iii) *If* $(N, Y) \cong (\mathbb{D}^2, \emptyset)$, $(\mathbb{D}^2, 0)$, $(\mathbb{S}^1 \times [0,1], \emptyset)$, $(\mathbb{M}, \emptyset)$, $(\mathbb{S}^2, 1pt)$, $(\mathbb{S}^2, 2pts)$, $(\mathbb{P}^2, 1pt)$, $(\mathbb{K}^2, \emptyset)$, *then* $\mathcal{H}_Y(N)_0 \simeq \mathbb{S}^1$.

Proof. In [12] the PL-homeomorphism groups of compact 2-manifolds was studied in the context of semisimplicial complex. However, using Corollary 2.1 and the results in [4], we can apply the arguments and results in [12, §3] to our setting.

(i) Let L be a small regular neighborhood of the union Y_1 of the nondegenerate components of Y and let $Y_0 = Y \setminus Y_1$. Since $\mathcal{H}_Y(N)_0$ deforms into $\mathcal{H}_{L \cup Y_0}(N)_0 \cong \mathcal{H}_{\mathrm{Fr}\, L \cup Y_0}(cl\,(N \setminus L))_0$, we may assume that $Y_1 \subset \partial N$. This case follows from [4] and [12, §3].

(ii) Let $\partial_+ N$ denote the union of the components of ∂N which meet A. Then $\mathcal{H}_{Y \cup A}(N)_0$ strongly deformation retracts onto $\mathcal{H}_{Y \cup \partial_+ N}(N)_0$, and the latter is contractible by the case (i). ☐

3. Relative isotopies on 2-manifolds

In Corollary 2.1 we have a principal bundle with a fiber $\mathcal{G} \equiv \mathcal{H}_X(M) \cap \mathcal{H}_K(M)_0$. In this section we will seek a sufficient condition which implies $\mathcal{G} = \mathcal{H}_X(M)_0$. Suppose M is a 2-manifold and N is a 2-submanifold of M. In [2] it is shown that (i) two homotopic essential simple closed curves in Int$M$ and two proper arcs homotopic rel ends in M are ambient isotopic rel ∂M, (ii) every homeomorphism $h : M \to M$ homotopic to id_M is ambient isotopic to id_M. Using these results or arguments we will show that if, in addition, $h|_N = id_N$ then h is isotopic to id_M rel N under some restrictions on disks, annuli and Möbius bands components (i.e. the pieces which admit global rotations). We denote the Möbius band, the torus and the Klein bottle by $\mathbb{M}$, $\mathbb{T}^2$ and $\mathbb{K}^2$ respectively. The symbol $\#X$ denotes the number of elements (or cardinal) of a set X.

Theorem 3.1. *Suppose M is a connected 2-manifold, N is a compact 2-submanifold of M and X is a subset of N such that*
(i) $M \neq \mathbb{T}^2, \mathbb{P}^2, \mathbb{K}^2$ *or* $X \neq \emptyset$.
(ii)(a) *if H is a disk component of N, then* $\#(H \cap X) \geq 2$,
(b) *if H is an annulus or Möbius band component of N, then* $H \cap X \neq \emptyset$,
(iii) (a) *if L is a disk component of $cl(M \setminus N)$, then* FrL *is a disjoint union of arcs or* $\#(L \cap X) \geq 2$,
(b) *if L is a Möbius band component of $cl(M \setminus N)$, then* FrL *is a disjoint union of arcs or* $L \cap X \neq \emptyset$.
If $h_t : M \to M$ is an isotopy rel X such that $h_0|_N = h_1|_N$, then there exists an isotopy $h'_t : M \to M$ rel N such that $h'_0 = h_0$, $h'_1 = h_1$ and $h'_t = h_t\ (0 \leq t \leq 1)$ on $M \setminus K$ for some compact subset K of M.

Corollary 3.1. *Under the same condition as in Theorem 3.1, we have* $\mathcal{H}_N(M) \cap \mathcal{H}_X(M)_0 = \mathcal{H}_N(M)_0$.

First we explain the meaning of the conditions (ii) and (iii) in Theorem 3.1. Suppose $h \in \mathcal{H}_N(M)$ and h is isotopic to id_M rel X. In order that h is isotopic to id_M rel N, it is necessary that h does not Dehn twist along the boundary circle of any disk, Möbius band and annulus component of N. This is ensured by the condition (ii) in Theorem 3.1 (Figure 1.*a*: $N = N_1 \cup N_2$, $cl(M \setminus N) = L$ and

$X = \{a, b\}$). However, this is not sufficient because a union of some components of N and $cl(M \setminus N)$ may form a disk, a Möbius band or an annulus. The condition (iii) in Theorem 3.1 is imposed to prevent Dehn twists around these pieces (Figure 1.*b*: $N = N_1 \cup N_2$, $cl(M \setminus N) = L \cup L_1 \cup L_2$ and $X = \{a_1, a_2, b_1, b_2\}$). This condition is too strong (we can replace X by $Y = \{a_1, b_1\}$), but it is simple and sufficient for our purpose.

FIGURE 1. h can not Dehn twist L.

Figure 1.a

Figure 1.b

We proceed to the verification of Theorem 3.1. We need some preliminary lemmas. Throughout this section we assume that M is a connected 2-manifold and N is a 2-submanifold of M. When G is a group and $S \subset G$, $\langle S \rangle$ denotes the subgroup of G generated by S.

We will use the fallowing facts from [2].

Fact 3.1. (0) ([2, Theorem 1.7]) If a simple closed curve C in M is null-homotopic, then it bounds a disk.
(1) ([2, Theorem 3.1]) Suppose α and β are proper arcs in M. If they are homotopic relative to end points, then they are ambient isotopic relative to ∂M.
(2) ([2, Theorem 4.2]) Let C be a simple closed curve in M, which does not bound a disk or a Möbius band. Let $\alpha \in \pi_1(M, *)$ be represented by a single circuit of C and let $\alpha = \beta^k$, $k \geq 0$. Then $\alpha = \beta$.
(3) ([2, Lemma 4.3]) (i) If $M \neq \mathbb{P}$, then $\pi_1(M)$ has no torsion elements.
(ii) Suppose $M \neq \mathbb{T}^2$, $\mathbb{K}^2$. If $\alpha, \beta \in \pi_1(M)$ and $\alpha\beta = \beta\alpha$, then $\alpha, \beta \in \langle \gamma \rangle$ for some $\gamma \in \pi_1(M)$.
(4) ([2, p101, lines 5 -10]) If $M \neq \mathbb{P}^2$ and every circle component of $\mathrm{Fr}N$ is essential in M, then the inclusion induces a monomorphism $\pi_1(N, x) \to \pi_1(M, x)$ for every $x \in N$.
(5) Suppose M is compact, X is a closed subset of ∂M, $X \neq \emptyset$ and $h : M \to M$ is a homeomorphism with $h|_X = id_X$.
(i) ([2, Theorem 3.4]) If $M = \mathbb{D}^2$ or $\mathbb{M}$ and $h|_{\partial M} : \partial M \to \partial M$ is orientation preserving, then h is isotopic to id_M rel X.
(ii) ([2, Proof of Theorem 6.3]) If $M \neq \mathbb{D}^2$ and h satisfies the following condition $(*)$, then h is isotopic to id_M rel X:
$(*)$ $h\ell \simeq \ell$ rel end points for every proper arc $\ell : [0, 1] \to M$ with $\ell(0), \ell(1) \in X$ (we allow that $\ell(0) = \ell(1)$ when X is a single point).

Comments. (4) Consider the universal covering $\pi : \tilde{M} \to M$. By Fact 3.1.(3-i) $\pi^{-1}(\mathrm{Fr}_M N)$ is a union of real lines, half rays and proper arcs. If $M = \mathbb{P}^2$, then $N = \mathbb{P}^2$.
(5-ii) M is a disk with k holes, ℓ handles (a handle = a torus with a hole) and m Möbius bands. The assertion is easily verified by the induction on $n = k + \ell + m$, using Fact 3.1.(1) and (5-i), together with the following remarks:

(a) When $\#X \geq 2$, we have $h\ell \simeq \ell$ rel. end points even if $\ell(0) = \ell(1)$.
(b) If h is (ambient) isotopic to h_1 rel. X, then h_1 also satisfies the condition $(*)$.
(c) Since $M \neq \mathbb{D}^2$, from the condition $(*)$ it follows that for every component C of ∂M, we have $h(C) = C$ and h preserves the orientation of C.
(d) Let $C_1, \cdots, C_p$ be the components of ∂M which meet X. Then h is isotopic rel. X to h_1 such that $h_1 = id$ on each C_i. Furthermore, h_1 satisfies $(*)$ for $\cup_i C_i$.

We also need the following remarks.

Fact 3.2. Suppose M is a connected 2-manifold and C is a circle component of ∂M. If either (i) $C \neq \partial M$ or (ii) M is noncompact, then C is a retract of M.

Comments. (ii) Take a half lay ℓ connecting C and ∞, and consider the regular neighborhood N of $C \cup \ell$. Since ∂N is a real line we can retract M onto N and then onto C.

Fact 3.3. Suppose M is a compact 2-manifold, $\{M_i\}$ is a finite collection of compact connected 2-manifolds such that $M = \cup_i M_i$ and $\mathrm{Int}M_i \cap \mathrm{Int}M_j = \emptyset$ $(i \neq j)$.
(i) If M is a disk, then some M_i is a disk.
(ii) If M is a Möbius band, then some M_i is a disk or a Möbius band.
(iii) If M is an annulus, then some M_i is a disk or an essential annulus in M. (If N is a disk with r holes in M $(r \geq 2)$, then there exists a disk $D \subset \mathrm{Int}M$ such that $D \cap N = \partial D \subset \partial N$.)

Lemma 3.1. *Suppose $M \neq \mathbb{K}^2$, C is a simple closed curve in M which does not bound a disk or a Möbius band in M, $x \in C$ and $\alpha \in \pi_1(M, x)$ is represented by C. If $\beta \in \pi_1(M, x)$ and $\beta^k = \alpha^\ell$ for some k, $\ell \in \mathbb{Z} \setminus \{0\}$, then $\beta \in \langle \alpha \rangle$.*

Proof. Attaching $\partial M \times [0, 1)$ to $\partial M \subset M$, we may assume that $\partial M = \emptyset$. Take a covering $p : (\tilde{M}, \tilde{x}) \to (M, x)$ such that $p_* \pi_1(\tilde{M}, \tilde{x}) = \langle \alpha, \beta \rangle \subset \pi_1(M, x)$.

If $\tilde{M}$ is noncompact, then by [2, Lemma 2.2] there exists a compact connected 2-submanifold N of $\tilde{M}$ such that $\tilde{x} \in N$ and the inclusion induces an isomorphism $\pi_1(N, \tilde{x}) \to \pi_1(\tilde{M}, \tilde{x})$. Since $\partial N \neq \emptyset$, it follows that $\pi_1(N, \tilde{x}) \cong \langle \alpha, \beta \rangle$ is a free group, so it is an infinite cyclic group $\langle \gamma \rangle$. By Fact.3.1.(2) $\gamma = \alpha^{\pm 1}$, so $\beta \in \langle \alpha \rangle$.

Suppose $\tilde{M}$ is compact. Since $\mathrm{rank} H_1(\tilde{M}) = 0$ or 1 and $\pi_1(\tilde{M}) \neq 1$, it follows that $\tilde{M} \cong \mathbb{P}^2$ or $\mathbb{K}^2$ and M is closed and nonorientable. If $\tilde{M} \cong \mathbb{K}^2$ so $\chi(\tilde{M}) = 0$, then $\chi(M) = 0$ and $M \cong \mathbb{K}^2$, a contradiction. Therefore, $\tilde{M} \cong \mathbb{P}^2$ and $\chi(\tilde{M}) = 1$, so $\chi(M) = 1$ and $M \cong \mathbb{P}^2$. We have $\pi_1(M) = \langle \alpha \rangle$. □

Note that if $M = \mathbb{K}^2$ and α, β are represented by the center circles of two Möbius bands, then $\alpha^2 = \beta^2$, but $\beta \notin \langle \alpha \rangle$.

Lemma 3.2. *Suppose C is a circle component of ∂M, $x \in C$ and $\alpha \in \pi_1(M,x)$ is represented by C. If $M \neq \mathbb{D}^2$, $\mathbb{M}$ or $\mathbb{S}^1 \times [0,1] \setminus A$ (A is a compact subset of $\mathbb{S}^1 \times \{1\}$), then there exists a $\gamma \in \pi_1(M,x)$ such that $\gamma\alpha^n \neq \alpha^n\gamma$ for any $n \in \mathbb{Z} \setminus \{0\}$.*

Proof. By the claim below we have a $\gamma \in \pi_1(M,x) \setminus \langle\alpha\rangle$. If $\gamma\alpha^n = \alpha^n\gamma$ for some $n \neq 0$, then by Fact.3.1.(3-ii) $\alpha^n, \gamma \in \langle\beta\rangle$ for some $\beta \in \pi_1(M,x)$ and $\alpha^n = \beta^k$ for some $k \in \mathbb{Z}$. Since $\alpha \neq 1$ and $M \neq \mathbb{P}^2$, by Fact.3.1.(3-i) $k \neq 0$. Hence by Lemma 3.1 $\beta \in \langle\alpha\rangle$ so $\gamma \in \langle\alpha\rangle$, a contradiction. $\square$

Claim. *Suppose M is a connected 2-manifold, C is a circle component of ∂M, $x \in C$ and $\alpha \in \pi_1(M,x)$ is represented by C. If $\pi_1(M,x) = \langle\alpha\rangle$, then $M \cong \mathbb{D}^2$ or $\mathbb{S}^1 \times [0,1] \setminus A$ for some compact subset A of $\mathbb{S}^1 \times \{1\}$.*

Proof. First we note that M does not contain any handles or Möbius bands. In fact if H is a handle or a Möbius band in M, then we can easily construct a retraction $r : M \to H$ which maps C homeomorphically onto ∂H (Fact 3.2), and we have the contradiction $\pi_1(H) = \langle r_*\alpha\rangle$. In particular, if M is compact then M is a disk or an annulus.

Suppose M is noncompact. It follows that ∂M contains no circle components other than C. In fact if C' is a circle in $\partial M \setminus C$, then we can join C and C' by a proper arc A in M and by Fact 3.2 we have a retraction $M \to C \cup A \cup C'$, a contradiction. We can write $M = \cup_{i=1}^{\infty} N_i$, where N_i is a compact connected 2-submanifold of M, $C \subset \mathrm{Int}_M N_1$, $N_i \subset \mathrm{Int}_M N_{i+1}$ and each component of $cl(M \setminus N_i)$ is noncompact. We will show that each $N = N_i$ is an annulus. This easily implies the conclusion.

Let $C_1, \cdots, C_m$ be the components of $\partial N \setminus C$. By the above remark $C_j \not\subset \partial M$, so C_j meets a component of $cl(M \setminus N)$. Let N' be a submanifold of N obtained by removing an open color of each C_j from N. It follows that $N' \cong N$, $\mathrm{Fr}N'$ is the union of circles C_j' associated with C_j's, each C_j' is contained in some component L_j of $cl(M \setminus N')$, $cl(M \setminus N') = \cup_j L_j$, and each L_j is noncompact. Since M contains no handles or Möbius bands (so no one point union of two circles), it follows that $L_j \cap L_{j'}' = \emptyset$ $(j \neq j')$ and $L_j \cap N' = C_j'$. By Fact 3.2 N' is a retract of M, so $\pi_1(N',x) = \langle\alpha\rangle$. This implies that $N \cong N'$ is an annulus. $\square$

The next lemma is a key point in the proof of Theorem 3.1. In [2, Lemma 6.1] the condition "the loop $h_t(x)$ is null-homotopic in M" is achieved by rotating x along C. However, this process does not keep the condition "isotopic rel N".

Lemma 3.3. *Suppose C is a circle component of $\mathrm{Fr}N$ which does not bound a disk or a Möbius band in M, $h : M \to M$ is a homeomorphism with $h|_N = id_N$ and $h_t : M \to M$ $(0 \leq t \leq 1)$ is a homotopy with $h_0 = h$, $h_1 = id_M$. If the following conditions are satisfied, then for any $x \in C$ the loop $m = \{h_t(x) : 0 \leq t \leq 1\}$ is null-homotopic in M :*
(i) *$M \neq \mathbb{T}^2$, $\mathbb{P}^2$, $\mathbb{K}^2$,*
(ii) *each circle component of $\mathrm{Fr}N$ is essential in M,*
(iii) *each component of $N \not\cong \mathbb{D}^2$, $\mathbb{M}$, $\mathbb{S}^1 \times [0,1] \setminus A$ ($A \subset \mathbb{S}^1 \times \{1\}$, compact).*

Proof. Let $\alpha = \{\ell\} \in \pi_1(M, x)$ be represented by C and let $\beta = \{m\} \in \pi_1(M, x)$. The homotopy $h_t\ell$ implies that $\alpha\beta = \beta\alpha$. Since $M \not\cong \mathbb{T}^2, \mathbb{K}^2$, by Fact.3.1.(3-ii) $\langle \alpha, \beta \rangle \subset \langle \delta \rangle$ for some $\delta \in \pi_1(M, x)$. Since C does not bound a disk or a Möbius band, by Fact 3.1.(2) $\delta = \alpha^{\pm 1}$ so $\beta = \alpha^k$ for some $k \in \mathbb{Z}$. Let $\alpha_1 = \{\ell\} \in \pi_1(N, x)$. By Lemma 3.2 there exists a $\gamma = \{n\} \in \pi_1(N, x)$ such that $\gamma\alpha_1^i \neq \alpha_1^i\gamma$ for any $i \in \mathbb{Z} \setminus \{0\}$ (Figure 2). The homotopy h_tn implies that $\gamma\beta = \beta\gamma$ in $\pi_1(M, x)$. Since $\pi_1(N, x) \to \pi_1(M, x)$ is monomorphic by Fact.3.1.(4), $\gamma\alpha_1^k = \alpha_1^k\gamma$ in $\pi_1(N, x)$ so that $k = 0$ and $\beta = 1$ in $\pi_1(M, x)$. $\square$

FIGURE 2. The loops ℓ, m and n in Lemma 3.3.

Figure 2

Lemma 3.4. *Suppose $N \neq \emptyset$, $cl(M \setminus N)$ is compact, each component of* FrN *is a circle, $h : M \to M$ is a homeomorphism such that $h|_N = id_N$ and h is homotopic to id_M. If the following conditions are satisfied, then h is isotopic to id_M rel N :*
(i) $M \neq \mathbb{T}^2$, $\mathbb{P}^2$, $\mathbb{K}^2$,
(ii) *each component C of* FrN *does not bound a disk or a Möbius band,*
(iii) *each component of $N \not\cong \mathbb{S}^1 \times [0, 1] \setminus A$ ($A \subset \mathbb{S}^1 \times \{1\}$, compact).*

If we assume that h is isotopic to id_M, then the condition (iii) is weakened to the condition:
(iii)$'$ each component of $N \not\cong \mathbb{S}^1 \times [0, 1]$, $\mathbb{S}^1 \times [0, 1)$.

Proof. Let $h_t : h \simeq id_M$ be any homotopy and let $L_1, \cdots, L_m$ be the components of $cl(M \setminus N)$. By Lemma 3.3 the loop $h_t(x) \simeq *$ in M for any $x \in \mathrm{Fr}N = \cup_j \mathrm{Fr}L_j$. We must find an isotopy $h|_{L_j} \simeq id_{L_j}$ rel FrL_j.

Let $f : [0, 1] \to L_j$ be any path with $f(0)$, $f(1) \in \mathrm{Fr}L_j$. The homotopy h_tf yields a contraction of the loop $hf \cdot h_t(f(1)) \cdot f^{-1} \cdot (h_t(f(0)))^{-1}$ in M. Since $h_t(f(0))$, $h_t(f(1)) \simeq *$, it follows that $hf \cdot f^{-1} \simeq *$ in M. Since $\pi_1(L_j) \to \pi_1(M)$ is monomorphic by Fact.3.1.(4), the loop $hf \cdot f^{-1} \simeq *$ in L_j, and the desired isotopy is obtained by Fact.3.1.(5-ii). $\square$

Figure 3 illustrates an original idea to prove Lemma 3.4 and Theorem 3.1: Consider the loop $m = n_1fn_2f^{-1}$ (f^{-1} is the inverse path of f). Any isotopy $h_t : id_M \simeq h$ rel $\{a, b\}$ induces a homotopy $h_tm : m \simeq n_1(hf)n_2(hf)^{-1}$ in $M \setminus \{a, b\}$. Modify the homotopy h_tm to simplify the intersection of the image of h_tm and FrN, and obtain a homotopy $F : \mathbb{S}^1 \times [0, 1] \to M \setminus \{a, b\}$ shown in Figure 3. The homotopies $F|_{A_1}$ in N_1 and $F|_{A_2}$ in $N_2 \setminus \{a, b\}$ imply that $k_i = 0$ ($i = 1, 2$), and the homotopy $F|_{B_1}$ in L implies that $f \simeq hf$ rel end points in L as required.

Proof of Theorem 3.1. We can assume that X is a finite set, since there exists a finite subset Y of X such that (M, N, Y) satisfies the conditions (i) - (iii) in Theorem 3.1. Replacing h_t by $h_1^{-1}h_t$, we may assume that $h_1 = id_M$.

FIGURE 3.

Figure 3

(I) The case where M is compact : Let $N_1, \cdots, N_n$ be the components of N and $L = cl(M \setminus N)$. Let $K_1, \cdots, K_p$ be the components of L which are disks or Möbius bands and let $L_1, \cdots, L_q$ be the remaining components. For each j we can write

$$\partial L_j = \left(\cup_{i=1}^{k(j)} A_i^j\right) \cup \left(\cup_{i=1}^{\ell(j)} B_i^j\right) \cup \left(\cup_{i=1}^{m(j)} C_i^j\right),$$

where A_i^j's are the circle components of $\mathrm{Fr}L_j$, B_i^j's are the components of ∂L_j which contain some arc components of $\mathrm{Fr}L_j$ and C_i^j's are the remaining components of ∂L_j. We choose disjoint collars E_i^j of A_i^j and F_i^j of B_i^j in L_j and set $\hat{A}_i^j = \partial E_i^j \setminus A_i^j$, $\hat{B}_i^j = \partial F_i^j \setminus B_i^j$ and

$$L_j' = cl(L_j \setminus (\cup_{i=1}^{k(j)} E_i^j) \cup (\cup_{i=1}^{\ell(j)} F_i^j)), \quad N' = N \cup (\cup_{k=1}^{p} K_k) \cup \left(\cup_{j=1}^{q} \left[\left(\cup_{i=1}^{k(j)} E_i^j\right) \cup \left(\cup_{i=1}^{\ell(j)} F_i^j\right)\right]\right).$$

Note that $\mathrm{Fr}N' = \cup_{j=1}^{q} \left[\left(\cup_{i=1}^{k(j)} \hat{A}_i^j\right) \cup \left(\cup_{i=1}^{\ell(j)} \hat{B}_i^j\right)\right] \subset \mathrm{Int}M$. Since $\mathcal{H}_\partial(\mathbb{D}) \simeq \mathcal{H}_\partial(\mathbb{M}) \simeq *$ by Fact 3.1.(5-i), we can isotope h_0 rel N to an $h' \in \mathcal{H}_{N'}(M)$.

By the construction (M, N', X, h') satisfies the following conditions:

(1) N' is a 2-submanifold of M, every component of $\mathrm{Fr}N'$ is a circle and $X \subset \mathrm{Int}_M N'$.

(2) $h'|_{N'} = id_{N'}$ and h' is isotopic to id_M rel X.

(3) Suppose C is a component of $\mathrm{Fr}N'$. If C bounds a disk D then $\#(D \cap X) \geq 2$, and if C bounds a Möbius band E then $E \cap X \neq \emptyset$.

(4) If H is an annulus component of N' then $H \cap X \neq \emptyset$.

To see (3) first note that M is the union of compact 2-manifolds N_i's, E_i^j's, F_i^j's, K_k's and L_j''s, which have disjoint interiors. Suppose G is a compact connected 2-manifold in M with $\partial G \subset \mathrm{Fr}N'$. Since $G \subset \mathrm{Int}M$ and each F_i^j meets ∂M, it follows that G is the union of N_i's, E_i^j's, L_j''s and K_k's contained in G. Since E_i^j is an annulus and $L_j' \cong L_j$ is not a disk or a Möbius band, from Fact 3.3 it follows that (i) if G is a disk, then G contains a disk which is some N_i or K_k with $K_k \subset G \subset \mathrm{Int}\, M$, so $\#(G \cap X) \geq 2$, (ii) if G is a Möbius band, then G contains a disk or a Möbius band which is some N_i or K_k with $K_k \subset G \subset \mathrm{Int}\, M$, so $G \cap X \neq \emptyset$.

As for (4), H is the union of N_i's E_i^j's, F_i^j's and K_k's contained in H, and H contains at least one N_i, which is a disk with r holes. If $r \leq 1$ then by the assumption $N_i \cap X \neq \emptyset$. If $r \geq 2$ then we can find a disk D in $\mathrm{Int}H$ such that $D \cap N_i = \partial D \subset \partial N_i$ (Fact 3.3.(iii)). Since $D \subset \mathrm{Int}N' \subset \mathrm{Int}M$, D is a union of N_i's and K_k's and we can conclude that it coincides with some $K_k (\subset \mathrm{Int}M)$, which meets X. These imply (4).

It remains to show that h' is isotopic to id_M rel N' under the conditions (1) - (4). (i) When $X \subset \mathrm{Int}N'$, we can apply Lemma 3.4 to the triple $(M \setminus X, N' \setminus X, h'|_{M \setminus X})$. To verify the condition (iii) in Lemma 3.4, note that (a) each component of $N' \setminus X$ takes of the form $H \setminus X$ for some component of H of N', and, in particular, (b) if $H \setminus X \cong \mathbb{S}^1 \times [0, 1)$, then H is a disk and $\#H \cap X \geq 2$, a contradiction. Therefore $h'|_{M\setminus X}$ is isotopic to $id_{M\setminus X}$ rel $N'\setminus X$. Extending this isotopy over M by id_X we have the required isotopy $h'\simeq id_M$ rel N'. (ii) In the case where $X\not\subset \mathrm{Int}N'$, let $C=\partial N'\cap\partial M$ and consider $(\tilde{M}=M\cup_C C\times[0,1], \tilde{N}=N'\cup_C C\times[0,1], X, \tilde{h})$, where $\tilde{h}$ is the extension of h' by $id_{C\times[0,1]}$. Then (a) $X\subset\mathrm{Int}\tilde{N}$ and $(\tilde{M},\tilde{N},X,\tilde{h})$ satisfies (1) - (4), and (b) an isotopy of $\tilde{h}$ to $id_{\tilde{M}}$ rel $\tilde{N}$ restricts to an isotopy of h' to id_M rel N'. (Alternatively, we can modify the isotopy of h' to id_M rel X to an isotopy rel $X\cup V$, where V is a neighborhood of $X\cap\partial M$ in M. We can replace X so that $X\subset\mathrm{Int}N'$.) This completes the proof of the case (I).

(II) The case where M is noncompact: Choose a compact connected 2-submanifolds L_0 and L of M such that $h_t(N)\subset\mathrm{Int}_M L_0$ $(0\le t\le 1)$ and $L_0\subset\mathrm{Int}_M L$. Let $N_1=N\cup cl(L\setminus L_0)$. Since N_1 is a subpolyhedron of L with respect to some triangulation of L (cf. [2]), by Corollary 2.1 we have the principal bundle : $\mathcal{H}(L)_0\to\mathcal{E}(N_1,L)^*_0$. Let $f_t\in\mathcal{H}(L)_0$, $f_1=id_L$, be any lift (= extension) of the path $e_t\in\mathcal{E}(N_1,L)^*_0$ defined by $e_t|_N=h_t|_N$ and $e_t=id$ on $cl(L\setminus L_0)$.

We can apply the case (I) to (L,N_1,X_1,f_t), $X_1=X\cup cl(L\setminus L_0)$. For the condition (iii) in Theorem 3.1, when E is a component of $cl(L\setminus N_1)=cl(L_0\setminus N)$, (a) if $E\cap\mathrm{Fr}L_0=\emptyset$, then E is a component of $cl(M\setminus N)$ and (b) if $E\cap\mathrm{Fr}L_0\neq\emptyset$, then E contains a component of $\mathrm{Fr}L_0$ and $\mathrm{Fr}L_0\subset cl(L\setminus L_0)\subset X_1$ (it also follows that $\mathrm{Fr}_L E$ is not connected since $E\cap\mathrm{Fr}N\neq\emptyset$, so if E is a disk or a Möbius band, then $\mathrm{Fr}_L E$ is a disjoint union of arcs).

Therefore we have an isotopy $k_t:L\to L$ rel N_1 such that $k_0=f_0$, $k_1=id_L$. We can extend f_t and k_t to M by id. The required isotopy h'_t is defined by $h'_t=k_tf_t^{-1}h_t$. ☐

Proof of Corollary 3.1. Let $\mathcal{G}_1$ denote the unit path-component of a topological group $\mathcal{G}$. Theorem 3.1 implies $\mathcal{H}_N(M)\cap\mathcal{H}_X(M)_1=\mathcal{H}_N(M)_1$. When M is compact, from Fact 2.1 it follows that $\mathcal{H}_K(M)_0=\mathcal{H}_K(M)_1$ for any compact subpolyhedron K of M. Since X can be replaced by a finite subset Y of X as in the above proof, we have $\mathcal{H}_N(M)\cap\mathcal{H}_X(M)_0\subset\mathcal{H}_N(M)\cap\mathcal{H}_Y(M)_0=\mathcal{H}_N(M)_0$. The noncompact case follows from the same argument when we will show that $\mathcal{H}_K(M)_0$ is an ANR (Propositions 4.1,4.2) in the next section.

4. The homotopy types of the identity components of homeomorphism groups of noncompact 2-manifolds

In this final section we will prove Theorem 1.1 and Corollary 1.1. Below we assume that M is a *noncompact* connected PL 2-manifold and X is a compact subpolyhedron of M. We set $M_0=X$ and write as $M=\cup_{i=0}^{\infty}M_i$, where for each $i\ge 1$ (a) M_i is a nonempty compact connected PL 2-submanifold of M and $M_{i-1}\subset\mathrm{Int}_M M_i$, (b) for each component L of $cl\,(M\setminus M_i)$, L is noncompact and $L\cap M_{i+1}$ is connected and (c) $M_1\cap\partial M\neq\emptyset$ if $\partial M\neq\emptyset$. Taking a subsequence, we have the following cases :

(i) each M_i is a disk, (ii) each M_i is an annulus, (iii) each M_i is a Möbius band, and

(iv) each M_i is not a disk, an annulus or a Möbius band.

In (ii) the inclusion $M_i\subset M_{i+1}$ is essential, otherwise a boundary circle of M_i bounds a disk component in $cl\,(M\setminus M_i)$, and it contradicts the condition (b).

Lemma 4.1. *In the cases $(i) - (iii)$ it follows that*

(i) (a) $\partial M = \emptyset \Longrightarrow M \cong \mathbb{R}^2$,

(b) $\partial M \neq \emptyset \Longrightarrow M \cong \mathbb{D} \setminus A$, *where A is a nonempty 0-dimensional compact subset of $\partial\mathbb{D}$,*

(ii) (a) $\partial M = \emptyset \Longrightarrow M \cong \mathbb{S}^1 \times \mathbb{R}^1$,

(b) $\partial M \neq \emptyset \Longrightarrow$

$(b)_1$ $M \cong \mathbb{S}^1 \times [0, 1)$

$(b)_2$ $M \cong \mathbb{S}^1 \times [0, 1) \setminus A$, *where A is a nonempty 0-dimensional compact subset of $\mathbb{S}^1 \times \{0\}$,*

$(b)_3$ $M \cong \mathbb{S}^1 \times [0, 1] \setminus A$, *where A is a nonempty 0-dimensional compact subset of $\mathbb{S}^1 \times \{0, 1\}$,*

(iii) (a) $\partial M = \emptyset \Longrightarrow M \cong \mathbb{P}^2 \setminus 1pt$,

(b) $\partial M \neq \emptyset \Longrightarrow M \cong \mathbb{M} \setminus A$, *where A is a nonempty 0-dimensional compact subset of $\partial\mathbb{M}$.*

In the case (ii)(b)$_3$ we may further assume that M_1 meets both $\mathbb{S}^1 \times \{0\}$ and $\mathbb{S}^1 \times \{1\}$.

We choose a metric d on M with $d \leq 1$ and metrize $\mathcal{H}_X(M)$ by the metric ρ defined by

$$\rho(f, g) = \sum_{i=1}^{\infty} \frac{1}{2^i} \sup_{x \in M_i} d(f(x), g(x)).$$

We separate the following two cases:

(I) $(M, X) \cong (\mathbb{R}^2, \emptyset)$, $(\mathbb{R}^2, 1pt)$, $(\mathbb{S}^1 \times \mathbb{R}^1, \emptyset)$, $(\mathbb{S}^1 \times [0, 1), \emptyset)$, $(\mathbb{P}^2 \setminus 1pt, \emptyset)$.

(II) (M, X) is not Case (I).

Case (II): First we treat Case (II) and prove the following statements:

Proposition 4.1. *In Case* (II)*, we have* (1) $\mathcal{H}_X(M)_0 \simeq *$ *and* (2) $\mathcal{H}_X(M)_0$ *is an ANR.*

We use the following notation: For each $j \geq 1$ let $U_j = \mathrm{Int}_M M_j$ and $L_j = \mathrm{Fr}_M M_j$, and for each $j > i \geq k \geq 0$ let $\mathcal{H}_{k,j} = \mathcal{H}_{M_k \cup (M \setminus U_j)}(M)_0$, $\mathcal{U}^i_{k,j} = \mathcal{E}_{M_k}(M_i, U_j)^*_0$ and let $\pi^i_{k,j} : \mathcal{H}_{k,j} \to \mathcal{U}^i_{k,j}$ denote the restriction map, $\pi^i_{k,j}(h) = h|_{M_i}$.

Lemma 4.2. (1) $\mathcal{H}_{k,j} \cong \mathcal{H}_{M_k \cup L_j}(M_j)_0$ *is an AR .*

(2) *The map $\pi^i_{k,j} : \mathcal{H}_{k,j} \to \mathcal{U}^i_{k,j}$ is a principal bundle with the structure group $\mathcal{H}_{k,j} \cap \mathcal{H}_{M_i}(M) \cong \mathcal{G}^i_{k,j} \equiv \mathcal{H}_{M_k \cup L_j}(M_j)_0 \cap \mathcal{H}_{M_i \cup L_j}(M_j)$ (under the restriction map).*

(3) *$\mathcal{U}^i_{k,j}$ is an open subset of $\mathcal{E}_{M_k}(M_i, M)^*_0$, $cl\,\mathcal{U}^i_{k,j} \subset \mathcal{U}^i_{k,j+1}$ and $\mathcal{E}_{M_k}(M_i, M)^*_0 = \cup_{j>i} \mathcal{U}^i_{k,j}$.*

Proof. The statement (1) follows from Fact 2.1 and Lemma 2.1.(ii), and (2) follows from Corollary 2.1. For (3), note that $\mathcal{E}_{M_k}(M_i, M)^*_0$ is path connected (Proposition 2.1) and each $f \in \mathcal{E}_{M_k}(M_i, M)^*_0$ is isotopic to the inclusion $M_i \subset M$ in a compact subset of M. □

Lemma 4.3. *In Case* (II)*, for each $j > i > k \geq 0$,* (a) *$\mathcal{G}^i_{k,j}$ is an AR,* (b) *the restriction map $\pi^i_{k,j} : \mathcal{H}_{k,j} \to \mathcal{U}^i_{k,j}$ is a trivial bundle and* (c) *$\mathcal{U}^i_{k,j}$ is also an AR.*

Proof. Once we show that $(*)$ $\mathcal{G}^i_{k,j} = \mathcal{H}_{M_i \cup L_j}(M_j)_0$, then (a) the fiber $\mathcal{G}^i_{k,j}$ is an AR by Fact 2.1 and Lemma 2.1.(ii), so (b) the principal bundle has a global section and it is trivial and (c) follows from Lemma 4.2.(1). It remains to prove $(*)$.

(1) The cases (i)(a), (ii)(a), (iii)(a), (ii)(b)$_1$ and (iv) (under the condition (II)): We can apply Theorem 3.1 to $(\tilde{M_j} = M_j \cup_{L_j} L_j \times [0,1], \tilde{M_i} = M_i \cup_{L_j} L_j \times [0,1], \tilde{M_k} = M_k \cup_{L_j} L_j \times [0,1])$. We can verify the conditions (ii) and (iii) in Theorem 3.1 as follows: (ii) By the assumption $(M_i, X) \not\cong (\mathbb{D}, \emptyset)$, $(\mathbb{D}, 1pt)$, $(\mathbb{S}^1 \times [0,1], \emptyset)$, $(\mathbb{M}, \emptyset)$ for each $i \geq 1$. (iii) If H is a component of $cl_{\tilde{M_j}}(\tilde{M_j} \setminus \tilde{M_i}) = cl_M(M_j \setminus M_i)$, then H contains a component of L_j. (Also, H meets both M_i and L_j (if $H \cap L_j = \emptyset$ then H is a compact component of $cl(M \setminus M_i)$, a contradiction.), so $\mathrm{Fr}_{\tilde{M_j}} H$ is not connected. Hence if H is a disk or a Möbius band, then $\mathrm{Fr}_{\tilde{M_j}} H$ is a disjoint union of arcs.) By Corollary 3.1 (Compact case) it follows that $\mathcal{H}_{\tilde{M_k}}(\tilde{M_j})_0 \cap \mathcal{H}_{\tilde{M_i}}(\tilde{M_j}) = \mathcal{H}_{\tilde{M_i}}(\tilde{M_j})_0$ and this implies $(*)$.

(2) The cases (i)(b), (ii)(b)$_3$ and (iii)(b): Since $M_1 \cap \partial M \neq \emptyset$ and M_1 meets both $\mathbb{S}^1 \times \{0\}$ and $\mathbb{S}^1 \times \{1\}$ in the case (ii)(b)$_3$, it follows that $cl(M_j \setminus M_i)$ is a disjoint union of disks, thus $\mathcal{H}_{M_i \cup L_j}(M_j) = \mathcal{H}_{M_i \cup L_j}(M_j)_0$ by Fact 3.1.(5-i). This implies $(*)$.

(3) The remaining case (ii)(b)$_2$: It follows that (a) $cl(M_j \setminus M_i)$ is a disjoint union of disks D_k and an annulus H and (b) $D_k \cap M_i$ is an arc, $D_k \cap L_j$ is a disjoint union of arcs ($\neq \emptyset$) and $H \cap M_i$, $H \cap L_j$ are the boundary circles of H. Since $N = M_i \cup (\cup_k D_k)$ is an annulus and $N \cap L_j \neq \emptyset$, from Theorem 3.1 it follows that $\mathcal{H}_{M_k \cup L_j}(M_j)_0 \cap \mathcal{H}_{N \cup L_j}(M_j) = \mathcal{H}_{N \cup L_j}(M_j)_0$. Each $f \in \mathcal{G}^i_{k,j}$ is isotopic rel $M_i \cup L_j$ to $f' \in \mathcal{H}_{N \cup L_j}(M_j)$. Since f' is isotopic to id rel $M_k \cup L_j$, it follows that $f' \in \mathcal{H}_{N \cup L_j}(M_j)_0$ and so $f \in \mathcal{H}_{M_i \cup L_j}(M_j)_0$. This completes the proof. □

Lemma 4.4. *In Case* (II)*, for each* $i \geq k \geq 0$*,*
(a) $\mathcal{E}_{M_k}(M_i, M)^*_0$ *is an AR,*
(b) *the restriction map* $\pi : \mathcal{H}_{M_k}(M)_0 \longrightarrow \mathcal{E}_{M_k}(M_i, M)^*_0$ *is a trivial principal bundle with fiber* $\mathcal{H}_{M_i}(M)_0$*,*
(c) $\mathcal{H}_{M_k}(M)_0$ *strongly deformation retracts onto* $\mathcal{H}_{M_i}(M)_0$*.*

Proof. By Lemma 4.3.(c) each $\mathcal{U}^i_{k,j}$ $(j > i)$ is an AR. Thus by Fact 4.2.(3) $\mathcal{E}_{M_k}(M_i, M)^*_0$ is also an AR and it strongly deformation retracts onto the single point set $\{M_i \subset M\}$. Hence the principal bundle

$$\mathcal{G}^i_k \equiv \mathcal{H}_{M_k}(M)_0 \cap \mathcal{H}_{M_i}(M) \subset \mathcal{H}_{M_k}(M)_0 \longrightarrow \mathcal{E}_{M_k}(M_i, M)^*_0$$

is trivial and $\mathcal{H}_{M_k}(M)_0$ strongly deformation retracts onto the fiber $\mathcal{G}^i_k$. In particular, $\mathcal{G}^i_k$ is connected and $\mathcal{G}^i_k = \mathcal{H}_{M_i}(M)_0$. □

Proof of Proposition 4.1.(1). By Lemma 4.4.(c), for each $i \geq 0$ there exists a strong deformation retraction h^i_t $(0 \leq t \leq 1)$ of $\mathcal{H}_{M_i}(M)_0$ onto $\mathcal{H}_{M_{i+1}}(M)_0$. A strong deformation retraction h_t $(0 \leq t \leq \infty)$ of $\mathcal{H}_X(M)_0$ onto $\{id_M\}$ is defined as follows:

$$\begin{aligned} h_t(f) &= h^i_{t-i} h^{i-1}_1 \cdots h^0_1(f) \quad (f \in \mathcal{H}_X(M)_0,\ i \geq 0,\ i \leq t \leq i+1) \\ h_\infty(f) &= id_M. \end{aligned}$$

Since $\operatorname{diam} \mathcal{H}_{M_i}(M)_0 \leq 1/2^i \to 0$, the map $h : \mathcal{H}_X(M)_0 \times [0,\infty] \to \mathcal{H}_X(M)_0$ is continuous.

(In the cases (i), (ii) and (iii), the same conclusion follows from Lemma 2.1.(i)(ii) by taking the end compactification of M.) $\square$

For the proof of Proposition 4.1.(2), we will apply Hanner's criterion of ANRs:

Fact 4.1. ([6]) A metric space X is an ANR iff for any $\varepsilon > 0$ there is an ANR Y and maps $f : X \to Y$ and $g : Y \to X$ such that gf is ε-homotopic to id_X.

Proof of Proposition 4.1.(2). By Lemma 4.4.(b) and Proposition 4.1.(1) for each $i \geq 1$ we have the trivial principal bundle

$$\mathcal{H}_{M_i}(M)_0 \subset \mathcal{H}_X(M)_0 \xrightarrow{\pi} \mathcal{E}_X(M_i, M)_0^* \quad \text{with} \quad \mathcal{H}_{M_i}(M)_0 \simeq *.$$

It follows that π admits a section s, and the map $s\pi$ is fiber preserving homotopic to $id_{\mathcal{H}_X(M)_0}$ over $\mathcal{E}_X(M_i, M)_0^*$. Since each fiber of π has diam $\leq 1/2^i$, this homotopy is a $1/2^i$-homotopy. Since $\mathcal{E}_X(M_i, M)_0^*$ is an ANR (Proposition 2.1), by Fact 4.1 $\mathcal{H}_X(M)_0$ is also an ANR. $\square$

Case (I): The next statements follow from Lemma 2.1.(iii) and Fact 2.1 by taking the end compactification of M.

Proposition 4.2. *In Case* (I), *we have* (1) $\mathcal{H}_X(M)_0 \simeq \mathbb{S}^1$ *and* (2) $\mathcal{H}_X(M)_0$ *is an ANR.*

Theorem 1.1 follows from Propositions 4.1, 4.2, and Corollary 1.1 now follows from the following characterization of ℓ_2-manifold topological groups.

Fact 4.2. ([1]) A topological group is an ℓ_2-manifold iff it is a separable, non locally compact, completely metrizable ANR.

Proof of Corollary 1.1. Since M is locally compact and locally connected, $\mathcal{H}(M)$ is a topological group and $\mathcal{H}_X(M)$ is a closed subgroup of $\mathcal{H}(M)$. Since M is locally compact and second countable, $\mathcal{H}(M)$ is also second countable. A complete metric ρ on $\mathcal{H}(M)$ is defined by

$$\rho(f, g) = d_\infty(f, g) + d_\infty(f^{-1}, g^{-1}), \qquad d_\infty(f, g) = \sum_{n=1}^{\infty} \frac{1}{2^n} \sup_{x \in M_n} d(f(x), g(x))$$

for $f, g \in \mathcal{H}(M)$, where d is a complete metric on M with $d \leq 1$. Since $\mathcal{H}_X(M)_0 \cong \mathcal{H}_X(M)_0 \times s$ [3], $\mathcal{H}_X(M)_0$ is not locally compact. Finally, by Propositions 4.1, 4.2 $\mathcal{H}_X(M)_0$ is an ANR. This completes the proof. $\square$

References

[1] Dobrowolski, T. and Toruńczyk, H., Separable complete ANR's admitting a group structure are Hilbert manifolds, *Top. Appl.*, 12 (1981) 229 - 235.

[2] Epstein, D. B. A., Curves on 2-manifolds and isotopies, *Acta Math.*, 155 (1966) 83 - 107.

[3] Geoghegan, R., On spaces of homeomorphisms, embeddings, and functions I: *Topology*, 11 (1972) 159 - 177.

[4] Hamstrom, M. E., Homotopy groups of the space of homeomorphisms on a 2-manifold, *Illinois J. Math.*, 10 (1966) 563 - 573.

[5] ______ and Dyer E., Regular mappings and the space of homeomorphisms on a 2-manifold, *Duke Math. J.*, 25 (1958) 521 - 532.

[6] Hanner, O., Some theorems on absolute neighborhood retracts, *Ark. Mat.* 1 (1951) 389 - 408.
[7] Hu, S. T., Theory of Retracts, Wayne State Univ. Press, Detroit, 1965.
[8] Jakobsche, W., The space of homeomorphisms of a 2-dimensional polyhedron is an ℓ^2-manifold, *Bull. Acad. Polon. Sci. Sér. Sci. Math.,* 28 (1980) 71 - 75.
[9] Luke, R. and Mason, W. K., The space of homeomorphisms on a compact two - manifold is an absolute neighborhood retract, *Trans. Amer. Math. Soc.,* 164 (1972), 275 - 285.
[10] Mason, W. K., The space of all self-homeomorphisms of a 2-cell which fix the cell's boundary is an absolute retract, *Trans. Amer. Math. Soc.,* 161 (1971), 185 - 205.
[11] Rushing, T.B., *Topological Embeddings,* Academic Press, New York, 1973.
[12] Scott, G. P., The space of homeomorphisms of 2-manifold, *Topology,* 9 (1970) 97 - 109.
[13] Toruńczyk, H., Characterizing Hilbert space topology, *Fund. Math.,* 111 (1981) 247 - 262.
[14] Yagasaki, T., Spaces of embeddings of compact polyhedra into 2-manifolds, to appear in Top. Appl.

DEPARTMENT OF MATHEMATICS, KYOTO INSTITUTE OF TECHNOLOGY, MATSUGASAKI, SAKYOKU, KYOTO 606, JAPAN
E-mail address: `yagasaki@ipc.kit.ac.jp`